arXiv:1110.4077v2 [cs.CC] 12 Jan 2012

The parameterised complexity of list problems on graphs of bounded treewidth

Kitty Meeks and Alexander Scott
Mathematical Institute, University of Oxford, 24-29 St Giles, Oxford OX1 3LB, UK
`{meeks,scott}@maths.ox.ac.uk`

January 13, 2012

Abstract

We consider the parameterised complexity of several list problems on graphs, with parameter treewidth or pathwidth. In particular, we show that List Edge Chromatic Number and List Total Chromatic Number are fixed parameter tractable, parameterised by treewidth, whereas List Hamilton Path is W[1]-hard, even parameterised by pathwidth. These results resolve two open questions of Fellows, Fomin, Lokshtanov, Rosamond, Saurabh, Szeider and Thomassen (2011).

1 Introduction

Many graph problems that are know to be NP-hard in general are fixed parameter tractable when parameterised by the treewidth k of the graph, that is they can be solved in time $f(k) \cdot n^{O(1)}$ for some function f. Often there even exists a linear-time algorithm to solve the problem on graphs of fixed treewidth [2, 6, 24].

This is the case for a number of graph colouring problems. Although it is NP-hard to determine whether an arbitrary graph is 3-colourable, the chromatic number of a graph of treewidth at most k can be computed in linear time (Arnborg and Proskurowski [2]). Similarly, while it is NP-hard to determine the edge-chromatic number of cubic graphs (Holyer [15]) and more generally d-regular graphs for any $d \geq 3$ (Leven and Galil [20]), this problem can again be solved in linear time on graphs of bounded treewidth (Zhou, Nakano and Nishizeki [27]). The pattern is the same for the total colouring problem: this is NP-hard even for regular bipartite graphs (McDiarmid and Sánchez-Arroyo [22]) but there exists a linear-time algorithm to solve the problem on graphs of bounded treewidth (Isobe, Zhou and Nishizeki [16]).

However, list versions of such problems cannot always be solved so efficiently on graphs of bounded treewith. The well-known list variant of vertex colouring is clearly NP-hard in general (as it generalises the standard colouring problem), but can also be solved in polynomial time on graphs of fixed treewidth k, and even in linear time on such graphs if the number of colours is bounded (Jansen and Scheffler [17]). However, in contrast with the linearity result for CHROMATIC NUMBER, it has been shown that when the number of colours is unbounded LIST COLOURING is W[1]-hard and so not (up to certain complexity theoretic assumptions) fixed parameter tractable (Fellows, Fomin, Lokshtanov, Rosamond, Saurabh, Szeider and Thomassen [11]). The list variants of edge-colouring and total colouring are harder still on graphs of bounded treewidth: both problems are NP-hard on series-parallel graphs (Zhou, Matsuo and Nishizeki [28]), which have treewidth at most two, and list edge-colouring is also NP-hard on outerplanar graphs (Marx [21]), another class of graphs with treewidth at most two.

Perhaps surprisingly, however, it can be easier to determine the minimum length of lists required to guarantee the existence of a proper list colouring for a graph G than to determine whether G admits a proper colouring with a particular set of lists. Alongside the negative result about the complexity of LIST COLOURING, in [11] the authors also use Courcelle's theorem [7] to prove the following result.

Theorem 1.1 ([11]). *The* LIST CHROMATIC NUMBER *problem, parameterised by the treewidth bound t, is fixed-parameter tractable, and solvable in linear time for any fixed t.*

In Section 2 we show that the same pattern extends to the list chromatic number problems for edge-colouring and total colouring: although these problems are both NP-hard on graphs of treewidth two, it is possible to determine the list edge chromatic number and list total chromatic number of graphs of bounded treewidth in linear time, answering an open question from [11].

The situation is similar for the problem of determining whether a given graph contains a Hamilton path. HAMILTON PATH is known to be computationally difficult in general, remaining NP-hard when restricted to planar, cubic, 3-connected graphs (Garey, Johnson and Tarjan [13]) or bipartite graphs (Krishnamoorthy [19]), but can be solved in linear time on graphs of bounded treewidth (Arnborg and Proskurowski [2]). In Section 3 we consider a list variant of the problem, and show that it is unlikely to be fixed parameter tractable, even parameterised by pathwidth, answering another open question from [11].

In the remainder of this section we define formally the problems whose complexity we consider, and give some background on the treewidth bound and the theory of parameterised complexity.

1.1 Problems considered

A *proper edge-colouring* of a graph G is an assignment of colours to the edges of G so that no two incident edges receive the same colour. If a set $\mathcal{L} = \{L_e : e \in E(G)\}$ of lists of permitted colours is given, a *proper list edge-colouring* of $(G, \mathcal{L})$ is a proper edge-colouring of G in which each edge e receives a colour from its list L_e. The *list edge chromatic number* of G, $\mathrm{ch}'(G)$, is the smallest integer c such that, for any assignment of colour lists to the edges of G in which each list has length at least c, there exists a proper list edge-colouring of G. We define the following problem.

LIST EDGE CHROMATIC NUMBER
Input: A graph $G = (V, E)$.
Question: What is $\mathrm{ch}'(G)$?

A *proper total colouring* of a graph $G = (V, E)$ is an assignment of colours to the vertices and edges of G such that no two adjacent vertices or incident edges have the same colour, and no edge has the same colour as either of its endpoints. If a set $\mathcal{L} = \{L_x : x \in V \cup E\}$ of lists of permitted colours is given, a *proper list total colouring* of $(G, \mathcal{L})$ is a proper total colouring of G in which every vertex and edge receives a colour from its list. The *list total chromatic number* of G, $\mathrm{ch}_T(G)$, is the smallest integer c such that, for any assignment of colour lists to the vertices and edges of G in whcih each list has length at least c, there exists a proper list total colouring of G. We define the following problem.

List Total Chromatic Number

Input: A graph $G = (V, E)$.
Question: What is $\mathrm{ch}_T(G)$?

We also consider a list version of Hamilton Path, introduced in [11], where each vertex has a list of permissible positions on the path.

List Hamilton Path

Input: A graph $G = (V, E)$, and a set of lists $\mathcal{L} = \{L_v \subseteq \{1, \ldots, |V|\} : v \in V\}$ of permitted positions.
Question: Does there exist a path $P = P[1] \ldots P[|G|]$ in G such that, for $1 \leq i \leq |G|$, we have $i \in L_{P[i]}$?

1.2 Treewidth and Pathwidth

We consider the complexity of these problems restricted to graphs of bounded treewidth or pathwidth. Given a graph G, we say that $(T, \mathcal{D})$ is a *tree decomposition* of G if T is a tree and $\mathcal{D} = \{\mathcal{D}(t) : t \in T\}$ is a collection of non-empty subsets of $V(G)$ (or *bags*), indexed by the nodes of T, satisfying:

1. $V(G) = \bigcup_{t \in T} \mathcal{D}(t)$,
2. for every $e = uv \in E(G)$, there exists $t \in T$ such that $u, v \in \mathcal{D}(t)$,
3. for every $v \in V$, if $T(v)$ is defined to be the subgraph of T induced by nodes t with $v \in \mathcal{D}(t)$, then $T(v)$ is connected.

The *width* of the tree decomposition $(T, \mathcal{D})$ is defined to be $\max_{t \in T} |\mathcal{D}(t)| - 1$, and the *treewidth* of G is the minimum width over all tree decompositions of G. A *path decomposition* is a tree decomposition $(P, \mathcal{D})$ in which the indexing tree is a path, and the *pathwidth* of G is the minimum width over all path decompositions of G.

Given a tree decomposition $(T, \mathcal{D})$ of G, we assume that an arbitrary node $r \in V(T)$ is chosen to be the *root* of T, and define the *height*, $h(t)$ of any $t \in T$ to be the distance from r to t. For any $v \in V(G)$, we then define t_v to be the unique node t of minimum height such that $v \in \mathcal{D}(t)$ (i.e. t_v is the node of minimal height in the subtree $T(V)$).

We make use of the following bound on the order of a tree indexing a tree decomposition.

Lemma 1.2. *Let G be a graph of order n and treewidth k. Then there exists a width k tree decomposition $(T, \mathcal{D})$ for G with $|T| \leq n$.*

Proof. Let $(T_0, \mathcal{D}_0)$ be any width k tree decomposition of G. We will transform this decomposition into one with the required properties by repeatedly contracting edges of T'. For $i \geq 0$, if uv is an edge of T_i such that either $\mathcal{D}_i(u) \subseteq \mathcal{D}_i(v)$ or $\mathcal{D}_i(v) \subseteq \mathcal{D}_i(u)$, we obtain T_{i+1} by contracting the edge uv to a single node w, and set $\mathcal{D}_{i+1}(w) = \mathcal{D}_i(u) \cup \mathcal{D}_i(v)$ (note $|\mathcal{D}_{i+1}(w)| \leq k$), with all other bags of $\mathcal{D}_{i+1}$ the same as $\mathcal{D}_i$. If there is no such edge uv, we stop and set $(T, \mathcal{D}) = (T_i, \mathcal{D}_i)$.

We claim that the decomposition $(T, \mathcal{D})$, obtained when this process terminates, is as required. Observe that for every node $t \in T$ we must have $t = t_v$ for some $v \in V(G)$: this is trivially true for the root (any vertex in its bag will do), and if any other $t \in T$ is not equal to t_v for some $v \in V$ then all vertices in $\mathcal{D}(t)$ also belong to the bag indexed by the parent of t, and so we would have contracted this edge. Thus we have a surjective mapping from $V(G)$ to $V(T)$, implying that $|T| \leq n$, as required. □

We shall assume throughout that tree decompositions of this form are given.

Graphs having treewidth at most k can alternatively be characterised as *partial k-trees*, as in [2, 27]. It follows immediately from this equivalent definition that if a graph G of order n is a partial k-tree (i.e. it has treewidth at most k) then G has at most kn edges.

1.3 Parameterised complexity

When considering the class of problems which are solvable in polynomial time on graphs of treewidth at most k, we aim, as mentioned above, to distinguish those which can be solved in time $f(k) \cdot n^{O(1)}$. Problems in this subclass are said to be *fixed parameter tractable*, parameterised by treewidth. A standard method of showing that a problem does *not* belong to this subclass (and so the best known algorithm has running time $n^{f(k)}$ for some unbounded function f) is to prove that it is W[1]-hard – the analogue of proving a problem to be NP-hard – which implies, up to similar complexity theoretic assumptions, that the problem is not fixed parameter tractable. Background on the theory of parameterised complexity can be found in [9, 23].

Just as with proving that a problem is NP-hard, the primary method for showing the intractability of a parameterised problem is to give a reduction from another problem that is known to be W[1]-hard. If we know that the parameterised decision problem Π with parameter k is difficult and want to show that Π' is just as hard then, given any instance (I, k) of Π, we want to be able to construct an instance (I', k') of Π' such that (I', k') is a yes-instance for Π' if and only if (I, k) is a yes-instance for Π. The idea is to show in this way that if Π' is fixed parameter tractable (with parameter k') then Π must also be (with parameter k). Thus we need to show not only that (I', k') can be computed from (I, k) in time polynomial in $|I|$ and that $|I'|$ is bounded by a polynomial function of $|I|$ (as we would for an NP-hardness reduction), but also that k' is bounded by some function (not necessarily polynomial) of k.

One useful W[1]-hard problem, which we use for a reduction in Section 3, is MULTICOLOUR CLIQUE (shown to be W[1]-hard by Fellows, Hermelin, Rosamond and Vialette in [10]): given a graph G, properly coloured with k colours, the problem is to determine whether there exists a clique in G containing one vertex of each colour.

2 List Chromatic Number Problems

The main result of this section is the following theorem.

Theorem 2.1. LIST EDGE CHROMATIC NUMBER *and* LIST TOTAL CHROMATIC NUMBER *are fixed parameter tractable, parameterised by the treewidth bound k, and are solvable in linear time for any fixed k.*

The key technical tools we use to prove this result are Theorems 2.8 and 2.9, which determine the list edge chromatic number and list total chromatic number respectively for graphs with fixed treewidth and large maximal degree.

2.1 Background

We begin by recalling some existing results about the edge chromatic number and list edge chromatic number of a graph. It is easy to see that, for any graph G, we have

$$\Delta(G) \leq \chi'(G) \leq \mathrm{ch}'(G) \leq 2\Delta(G) - 1,$$

where $\chi'(G)$ denotes the edge chromatic number of G and $\Delta(G)$ is the maximum degree of G. For the edge chromatic number, we have the much stronger result of Vizing:

Theorem 2.2 ([25])**.** *$\chi'(G)$ is equal to either $\Delta(G)$ or $\Delta(G) + 1$.*

In order to give a linear time algorithm to solve Edge Chromatic Number on graphs of bounded treewidth, Zhou, Nakano and Nishizeki [27] prove that, for graphs of fixed treewidth k and maximum degree $\Delta \geq 2k$, the edge chromatic number must in fact be equal to Δ.

There is no direct analogue of Vizing's theorem for the list edge chromatic number. The *List (Edge) Colouring Conjecture* (discussed in [1, 5, 14]) asserts that $\chi'(G) = \mathrm{ch}'(G)$ for any graph G, and would immediately imply Vizing's conjecture (1976) that $\mathrm{ch}'(G) \leq \Delta(G)+1$. However, neither of these conjectures has been proved except for certain special classes of graphs, and the best general bound on the list edge chromatic number is due to Kahn.

Theorem 2.3 ([18])**.** *For any $\epsilon > 0$, if $\Delta(G)$ is sufficiently large,*

$$\mathrm{ch}'(G) \leq (1 + \epsilon)\Delta(G).$$

In Theorem 2.8, we show that for a graph G of bounded treewidth and large maximum degree,

$$\mathrm{ch}'(G) = \chi'(G) = \Delta(G),$$

proving a special case of the List (Edge) Colouring Conjecture. Using this result, the List Edge Chromatic Number problem on graphs of bounded treewidth can be reduced to the case in which the maximum degree of the graph is bounded.

We prove an analogous result for the list total chromatic number. Once again, there exist trivial bounds for the (list) total chromatic number of an arbitrary graph:

$$\Delta(G) + 1 \leq \chi_T(G) \leq \mathrm{ch}_T(G) \leq 2\Delta(G) + 1,$$

where χ_T denotes the total chromatic number of the graph. It is a long-standing but unproved conjecture (the *Total Colouring Conjecture* [3, 26]) that

$$\chi_T(G) \leq \Delta(G) + 2.$$

In Theorem 2.9 we prove that for a graph G of bounded treewidth and large maximum degree,

$$\mathrm{ch}_T(G) = \chi_T(G) = \Delta(G) + 1,$$

and so once again it suffices to solve the problem for graphs with bounded maximum degree.

Of course, there is a correspondence between these colouring problems and the vertex-colouring problems discussed above. For any graph $G = (V, E)$, the *line graph* $L(G)$ of G has vertex set E, and $e, f \in E$ are adjacent in $L(G)$ if and only if e and f are incident in G. Then solving (for example) List Edge Chromatic Number for the graph G is equivalent to solving List Chromatic Number for $L(G)$. Similarly, we define the *total graph* $T(G)$ of G to have vertex set $V \cup E$, and edge set $E \cup \{ef : e, f \in E \text{ and } e, f \text{ incident in } G\} \cup \{ve : v \text{ is an endpoint of } e\}$, and solving List Total Chromatic Number for G is then equivalent to solving List Chromatic Number for $T(G)$. However, as the treewidth of $L(G)$ or $T(G)$ can in general be arbitrarily large even when G itself has small treewidth, results about the parameterised complexity of vertex colouring problems do not immediately transfer to the edge and total colouring cases.

If the maximum degree of G is bounded, however, the following result (proved in [8]) tells us that the treewidth of $L(G)$ is bounded by a constant multiple of that of G.

Lemma 2.4. *Let G be a graph of treewidth at most k, and maximum degree at most Δ. Then $L(G)$ has treewidth at most $(k+1)\Delta$.*

A similar result holds for the treewidth of $T(G)$.

Lemma 2.5. *Let G be a graph of treewidth at most k, and maximum degree at most Δ. Then $T(G)$ has treewidth at most $(k+1)(\Delta+1)$.*

Proof. If $(T, \mathcal{D})$ is a width k tree decomposition for G, it is easy to verify that $(T, \mathcal{D}')$, where $\mathcal{D}'(t) = \mathcal{D}(t) \cup \{uv \in E : \{u, v\} \cap \mathcal{D}(t) \neq \emptyset\}$, is a tree decomposition of $T(G)$ of width at most $(k+1)(\Delta+1)$. □

We will need two further results for our proofs. First, a theorem of Galvin concerning the list edge chromatic number of bipartite graphs:

Theorem 2.6 ([12]). *Let G be a bipartite graph. Then*

$$\mathrm{ch}'(G) = \chi'(G) = \Delta(G).$$

Finally, we make use of an algorithm of Bodlaender:

Theorem 2.7 ([4]). *For all $k \in \mathbb{N}$, there exists a linear-time algorithm that tests whether a given graph $G = (V, E)$ has treewidth at most k and, if so, outputs a tree-decomposition of G with treewidth at most k.*

2.2 Results and Proofs

In this section we prove our two technical results about the list edge chromatic number and list total chromatic number of graphs with bounded treewidth and large maximum degree, and then give fixed parameter algorithms to solve LIST EDGE CHROMATIC NUMBER and LIST TOTAL CHROMATIC NUMBER on graphs of bounded treewidth.

Theorem 2.8. *Let G be a graph with treewidth at most k and $\Delta(G) \geq (k+2)2^{k+2}$. Then $\mathrm{ch}'(G) = \Delta(G)$.*

Proof. As $\Delta \leq \mathrm{ch}'(G)$, it is sufficient to prove that, if G is a graph of treewidth at most k, we have $\mathrm{ch}'(G) \leq \max\{\Delta(G), (k+2)2^{k+2}\}$. Suppose this does not hold, and let G be a counterexample with as few edges as possible, so there exists a set $\mathcal{L} = (L_e)_{e \in E(G)}$ of colour-lists, all of length $\Delta_0 = \max\{\Delta(G), (k+2)2^{k+2}\}$, such that there is no proper list edge-colouring of $(G, \mathcal{L})$. We may assume without loss of generality that G contains no isolated vertices and so, by edge-minimality of G, we must have $\mathrm{ch}'(G') \leq \max\{\Delta(G'), (k+2)2^{k+2}\} \leq \Delta_0$ for any proper subgraph G' of G.

We may assume that every edge $e \in E(G)$ is incident with at least Δ_0 others: if e is incident with fewer than Δ_0 other edges then we can extend any proper list edge-colouring of $(G - e, \mathcal{L} \setminus \{L_e\})$ to a proper list edge-colouring of $(G, \mathcal{L})$. We will show that, under this assumption, there must exist a nonempty set of vertices U such that any proper list edge-colouring of $(G \setminus U, (L_e)_{e \in E(G \setminus U)})$ can be extended to a proper list edge-colouring of

$(G, \mathcal{L})$, contradicting the choice of G as an edge-minimal counterexample (as the fact there are no isolated vertices in G means $e(G \setminus U) < e(G)$).

Let us define $L \subset V(G)$ to be the set of vertices of degree at least $\Delta_0/2$, and note that every edge is incident with at least one vertex from L (as otherwise it can be incident with only $\Delta_0 - 1$ other edges), so $S = V(G) \setminus L$ is an independent set. Fix a width k tree decomposition $(T, \mathcal{D})$ of G, and choose $v \in L$ such that $h(t_v) = \max_{x \in L} h(t_x)$. We then set T' to be the subtree of T rooted at t_v, that is the subgraph of T induced by nodes u such that the path from u to the root contains t_v.

Set $X \subset V(G)$ to be $\bigcup_{t \in T'} \mathcal{D}(t)$, and $X' = X \setminus \mathcal{D}(t_v)$. We then make the following observations.

1. $L \cap X \subseteq \mathcal{D}(t_v)$: if any vertex $z \in L$ appears in a bag indexed by T' but does not appear in $\mathcal{D}(t_v)$, we must have $h(t_z) > h(t_v)$, contradicting the choice of v. This implies immediately that $X' \subseteq S$.

2. $\Gamma(X') \subseteq \mathcal{D}(t_v)$: no vertex from X' can appear in a bag of the decomposition not indexed by T', so clearly we have $\Gamma(X') \subseteq X$; but also, as $X' \subset S$, we have $\Gamma(X') \subseteq L$ and so we see $\Gamma(X') \subseteq L \cap X \subseteq \mathcal{D}(t_v)$.

3. $|X'| \geq \Delta_0/2 - k$: since v appears only in bags indexed by T', we have $\Gamma(v) \subseteq X \setminus \{v\}$, implying $|X| - 1 \geq d(v) \geq \Delta_0/2$, and we know $|X| - |X'| = |\mathcal{D}(t_v)| \leq k+1$, so we have $|X'| \geq |X| - k - 1 \geq \Delta_0/2 - k$.

As the neighbourhood of any vertex $x \in X'$ is contained in $\mathcal{D}(t_v)$, there are at most 2^{k+1} possibilities for the neighbourhood of such a vertex. Therefore there must exist some subset $U \subseteq X$ such that every vertex in U has the same neighbourhood, and

$$\begin{aligned}
|U| &\geq \frac{|X'|}{2^{k+1}} \\
&\geq \frac{\Delta_0/2 - k}{2^{k+1}} \\
&\geq \frac{(k+2)2^{k+1} - k}{2^{k+1}} \\
&\geq k+1 \\
&\geq |\Gamma(U)|.
\end{aligned}$$

Now let ϕ be a proper edge-colouring of $(G \setminus U, (L_e)_{e \in E(G \setminus U)})$. If we can extend ϕ to a proper edge-colouring of G in which every edge incident with U

also receives a colour from its list, then we have a proper list edge-colouring of $(G, \mathcal{L})$, giving the required contradiction.

Set $W = \Gamma(U)$, say $W = \{w_1, \ldots, w_r\}$ (where $r \leq |U|$), and let H be the complete bipartite subgraph of G induced by $U \cup W$. Suppose, for $1 \leq i \leq r$, that F_i is the set of colours already used by ϕ on edges incident with w_i, and for each $u \in U$ define the list L'_{uw_i} to be $L_{uw_i} \setminus F_i$. If we can properly colour the edges of H in such a way that each edge $e \in E(H)$ is given a colour from L'_e, then we can extend ϕ as required.

Observe that, for each i, $|F_i| \leq \Delta(G) - |U|$, and so we have $|L_{uw_i}| \geq \Delta_0 - \Delta(G) + |U| \geq |U|$. But as H is bipartite, with maximum degree $|U|$, we have (by Theorem 2.6)

$$\mathrm{ch}'(H) = \Delta(H) = |U| \ .$$

Therefore, as each list L'_e contains at least $|U|$ colours, there exists a proper edge colouring of H in which every edge e receives a colour from its list L'_e, completing the proof. $\square$

We use a very similar argument to prove the following result about the list total chromatic number.

Theorem 2.9. *Let G be a graph with treewidth at most k and $\Delta(G) \geq (k+2)2^{k+2}$. Then $\mathrm{ch}_T(G) = \Delta(G) + 1$.*

Proof. As $\mathrm{ch}_T(G) \geq \Delta(G) + 1$, it suffices to prove that, if G is a graph of treewidth at most k, we have $\mathrm{ch}_T(G) \leq \Delta_0 + 1 = \max\{\Delta(G), (k+2)2^{k+2}\} + 1$. Again, we prove the result by contradiction, so suppose this does not hold, and let G be a counterexample with as few edges as possible, so there exists a set $\mathcal{L} = (L_x)_{x \in V(G) \cup E(G)}$ of colour-lists, all of length $\Delta_0 + 1$, such that there is no proper list total colouring of $(G, \mathcal{L})$. We may assume without loss of generality that G contains no isolated vertices and so, by edge-minimality of G, we must have $\mathrm{ch}_T(G') \leq \max\{\Delta(G'), (k+2)2^{k+2}\} + 1 \leq \Delta_0 + 1$ for any proper subgraph G' of G.

We may assume that every edge $e \in E(G)$ is incident with at least $\Delta_0 - 1$ others: if e is incident with fewer than $\Delta_0 - 1$ other edges then at most Δ_0 colours can appear on vertices or edges incident with e, and so any proper list total colouring of $(G - e, \mathcal{L} \setminus \{L_e\})$ can be extended to a proper list total colouring of $(G, \mathcal{L})$. Thus, if we define $L' \subset V(G)$ to be the set of vertices of degree at least $(\Delta_0 - 1)/2$, every edge is incident with at least one vertex from L', and $S' = V(G) \setminus L'$ is an independent set.

Exactly as in the proof of Theorem 2.8, we can find a subset $U \subset S'$ such that all vertices of U have the same neighbourhood of size at most $k+1$, and

$$\begin{aligned} |U| &\geq \frac{(\Delta_0 - 1)/2 - k}{2^{k+1}} \\ &\geq \frac{((k+2)2^{k+2} - 1)/2 - k}{2^{k+1}} \\ &> k+1 \\ &\geq |\Gamma(U)|. \end{aligned}$$

As $G \setminus U$ is a proper subgraph of G, there exists a proper list total colouring ϕ of $(G \setminus U, \mathcal{L}')$ where $\mathcal{L}' = (L_x)_{x \in E(G\setminus U) \cup V(G \setminus U)}$. If we can extend ϕ to a proper total colouring of G in which every vertex from U and every edge incident with U also receives a colour from its list, then we have a proper list total colouring of $(G, \mathcal{L})$, giving the desired contradiction.

As in the proof of Theorem 2.8, set $W = \Gamma(U)$, say $W = \{w_1, \ldots, w_r\}$ (where $r \leq |U|$), and let H be the complete bipartite subgraph of G induced by $U \cup W$. Suppose, for $1 \leq i \leq r$, that F_i is the set of colours already used by ϕ on w_i and the edges incident with w_i, and for each $u \in U$ define L'_{uw_i} to be $L_{uw_i} \setminus F_i$.

Observe that, for each i, $|F_i| \leq 1 + \Delta(G) - |U|$, and so we have

$$|L_{uw_i}| \geq \Delta_0 + 1 - 1 - \Delta(G) + |U| \geq |U|.$$

As H is bipartite, with maximum degree $|U|$, we have (by Theorem 2.6) $\mathrm{ch}'(H) = \Delta(H) = |U|$. Therefore, as each list L'_e contains at least $|U|$ colours that are not already used by ϕ on edges or vertices incident with e, we can extend ϕ to a proper list colouring ϕ' including the edges incident with U. If we can then colour the vertices of U in such a way that no $u \in U$ receives a colour used by ϕ' on an adjacent vertex or incident edge, we can indeed extend ϕ to a proper list total colouring of G.

Note that every $u \in U$ has degree at most $k+1$, and so is adjacent to or incident with at most $2(k+1)$ vertices and edges of G. Thus, as $|L_u| \geq \Delta_0 + 1 > 2(k+1)$, there is at least one colour in L_u that is not used by ϕ' on a vertex adjacent to u or on an edge incident with u, and we can extend ϕ' to the vertices of U greedily. This gives a proper list total colouring of $(G, \mathcal{L})$, completing the proof. $\square$

Using these theorems, we can now prove our main complexity result.

Proof of Theorem 2.1. Let G be a graph on n vertices, with treewidth at most k, and set $f(k) = (k+2)2^{k+2}$. We can check in time $O(f(k)n)$ whether $\Delta(G) \geq f(k)$, and if this is the case then, by Theorem 2.8 or Theorem 2.9, we know the exact value of $\mathrm{ch}'(G)$ or $\mathrm{ch}_T(G)$ respectively. Thus it suffices to solve both problems in the case that $\Delta(G) < f(k)$.

This is exactly the same as solving List Chromatic Number on $L(G)$ or $T(G)$, when $\Delta(G) < f(k)$. But in this case, by Lemmas 2.4 and 2.5, $L(G)$ and $T(G)$ have bounded treewidth. Note that both graphs can be computed from G in time

$$\begin{aligned} O(e(L(G)) + e(T(G))) &= O((k+1)(f(k)+1)(|T(G)| + |L(G)|) \\ &= O(k^2 f(k)|G|), \end{aligned}$$

and so we can then use Bodlaender's algorithm (Theorem 2.7) to find a tree decomposition of $L(G)$ or $T(G)$, of width at most $(k+1)(f(k)+1)$, in time $O(|G|)$ for any fixed k. By Theorem 1.1, given this decomposition, we can compute the list chromatic number of $L(G)$ or $T(G)$, and hence the list edge chromatic number or list total chromatic number of G, in linear time. ☐

Our proof also implies a polynomial-time algorithm to compute a proper list edge-colouring of any graph G of fixed treewidth and large maximum degree, provided every $L \in \mathcal{L}$ has length at least $\Delta(G)$. The same method can also be used to compute a list total colouring of such a graph, provided every list has length at least $\Delta(G) + 1$.

Theorem 2.10. *Let G be a graph of order n and treewidth k, with maximum degree at least $(k+2)2^{k+3}$, and let $\mathcal{L} = (L_e)_{e \in E(G)}$ be a set of colour-lists such that $|L_e| \geq \Delta(G)$ for all $e \in E(G)$. Then, for fixed k, we can compute a proper list edge-colouring of $(G, \mathcal{L})$ in time $O(n^2)$.*

Proof. The idea of the algorithm is to delete repeatedly edges or sets of vertices, as in the proof of Theorem 2.8, until the degree of the remaining graph is less than $(k+2)2^{k+2} < \Delta(G)/2$, so the edges of this graph can be list-coloured greedily. Edges and vertices are then reinserted and coloured to extend this colouring to the edges of G. Unlike in the proof of Theorem 2.8, where we identified a set of *at least* $k+1$ vertices to delete, in this algorithm we always delete a set of *exactly* $k+1$ vertices.

We begin with some preprocessing. For each vertex, we construct a list of its neighbours, and we also construct an n-element array storing the degree of each vertex. For vertices of degree at most $k+1$, the list of neighbours is sorted into order, and the list of vertices of degree at most $k+1$ is also sorted, in order of neighbourhoods (so that vertices with the same neighbourhood occur consecutively). All this can be done in time $O(n^2)$.

Note that each time we delete an edge, we can update this information in time $O(n)$: we update the neighbour lists of the edge's endpoints and decrement their degrees, and if one or both of the endpoints now has degree at most $k+1$ its neighbours are sorted and it is inserted into the correct position in the list of small-degree vertices. From the point of view of updating information, deleting a set of $k+1$ vertices can be regarded as a series of edge-deletions, each performed in time $O(n)$. As there are $O(n)$ edges in total, this means we can perform all updates after deletions in time $O(n^2)$.

Given the array of degrees of all vertices, it is straightforward to identify in time $O(n)$ an edge incident with fewer than $\Delta(G)$ others, if such an edge exists. If there is no such edge, we know from the proof of Theorem 2.8 that (provided the maximum degree of the graph is still at least $(k+2)2^{k+2}$) there exists a set of $k+1$ vertices with a common neighbourhood of size at most $k+1$. As the vertices of degree at most $k+1$ are sorted by their neighbourhoods, it is also possible to identify such a set of vertices in linear time. Thus at each step we are able to identify the edge or set of vertices to delete in time $O(n)$, and so all deletions (and subsequent updating) can be performed in time $O(n^2)$.

It therefore remains to show that we can also perform the reinsertions in time $O(n^2)$. When we reinsert an edge, we can simply colour it with the first available colour from its colour-list, taking time $O(n)$. When reinserting a set of $k+1$ vertices, we need to colour up to $(k+1)^2$ edges which form a complete bipartite subgraph. Recall from the proof of Theorem 2.8 that every such edge still has at least $k+1$ available colours from its list (i.e. colours that have not already been used on incident edges), and that there exists a proper list edge-colouring of the bipartite graph if every edge has a list of exactly $k+1$ permitted colours. For each of the edges, we can compute in time $O(kn)$ a list of $k+1$ colours from its colour-list which have not yet been used on incident edges. We can then check in constant time all possible colourings of the $(k+1)^2$ edges in which each receives one of the $k+1$ colours from its list, to find a proper colouring of this bipartite graph, which is guaranteed to extend the list edge-colouring as required.

Thus we can perform all reinsertions in time $O(n^2)$, completing the proof of the theorem. $\square$

3 List Hamilton Path

Given a graph G and a set of lists $\mathcal{L} = \{L_v \subseteq \{1, \ldots, |G|\} : v \in V(G)\}$ of permitted positions, we say a path $P = P[1] \ldots P[|G|]$ in G is a *valid* Hamilton path if $i \in L_{P[i]}$ for every i.

We prove the following result.

Theorem 3.1. List Hamilton Path, *parameterised by pathwidth, is W[1]-hard.*

We prove the theorem by means of a reduction from Multicolour Clique. Suppose G is the k-coloured graph in an instance of Multicolour Clique: we may assume without loss of generality that all k vertex classes have the same size, and also that the number of edges between each pair of vertex classes is the same (as adding isolated edges and vertices does not change the existence or otherwise of a multicolour clique). Let the vertex classes be $V_1, \ldots, V_k$, where V_i contains vertices $V_i[1], \ldots, V_i[p]$, and assume that there are q edges between each pair of vertex classes. We describe the construction of a graph H of pathwidth at most $5k$, and a set of lists $\mathcal{L} = (L_v)_{v \in V(H)}$ such that there is a valid Hamilton path in $(H, \mathcal{L})$ if and only if G contains a multicolour clique. Lemmas 3.4, 3.5 and 3.6 will demonstrate that $(H, \mathcal{L})$ does indeed have the required properties.

Our construction consists of $k+1$ paths, with some additional edges linking them: paths $P_1, \ldots, P_k$ correspond to the vertex classes $V_1, \ldots, V_k$, and an additional path Q is used to connect $P_1, \ldots, P_k$. Each path P_i has $2n^2$ vertices (so $|P_i| \geq 4q\binom{k}{2} = 4e(G)$), and we denote the j^{th} vertex of P_i by $P_i[j-1]$. The path $Q = Q_1 \ldots Q_k$ is the concatenation of k subpaths $Q_1, \ldots, Q_k$, each containing $n^2(n-2)$ vertices and, for $1 \leq i \leq k$, every vertex of Q_i is adjacent to both $P_i[0]$ and $P_i[2n^2 - 1]$.

In addition, we have a number of edge-gadgets, consisting of edges between pairs of the paths $P_1, \ldots, P_k$. Suppose $E(G) = \{e_0, \ldots, e_{m-1}\}$. Then, for each edge e_r between V_i and V_j, we have an edge-gadget $G(e_r)$, involving the r^{th} group of four vertices in P_i and the corresponding group of vertices from P_j. $G(e_r)$ has edges $P_i[4r]P_j[4r+1]$, $P_j[4r]P_i[4r+1]$, $P_i[4r+1]P_j[4r+3]$ and $P_i[4r+2]P_j[4r+2]$, as illustrated in Figure 1.

This completes the construction of the graph H. We now define the list of permitted positions for each vertex. For $1 \leq i \leq k$, we set

$$L_{P_i[0]} = \{(i-1)n^3 + 2\alpha n^2 : 1 \leq \alpha \leq p\},$$



Figure 1: The edge-gadget $G(e_r)$

and

$$L_{P_i[2n^2-1]} = \{(i-1)n^3 + 2\alpha n^2 + (2n^2-1) + k + 1 - 2i : 1 \leq \alpha \leq p\}.$$

We further define the list

$$L(i,j) = \{(i-1)n^3 + 2\alpha n^2 + j + \beta : 1 \leq \alpha \leq p, -(k-1) \leq \beta \leq k-1\}.$$

For every internal vertex $P_i[j]$, the list $L_{P_i[j]}$ will contain $L(i,j)$. In most cases we in fact set $L_{P_i[j]} = L(i,j)$, the only exceptions being three vertices in each edge-gadget $G(e)$: if $e = V_i[r]V_j[s]$ and the vertices in $G(e)$ are $P_i[l], \ldots, P_i[l+3], P_j[l], \ldots, P_j[l+3]$, then the list for $P_i[l+1]$ will additionally contain positions

$$\{(j-1)n^3 + 2sn^2 + (l+1) + \beta : -(k-1) \leq \beta \leq k-1\},$$

while the lists for $P_j[l+1]$ and $P_j[l+2]$ also contain

$$\{(i-1)n^3 + 2rn^2 + (l+1) + \beta : -(k-1) \leq \beta \leq k-1\}$$

and

$$\{(i-1)n^3 + 2rn^2 + (l+2) + \beta : -(k-1) \leq \beta \leq k-1\}$$

respectively. We place no restriction on the positions that vertices from Q can take in a valid Hamilton path.

The idea is that any valid Hamilton path must (with the exception of a few vertices belonging to the edge-gadgets) traverse $P_1, \ldots, P_k$ in that order, using sections of Q before and after each P_i to connect the paths. In this construction, our choice of position for $P_i[0]$ corresponds to a choice of vertex from V_i: if $P_i[0]$ takes position $(i-1)n^3 + 2rn^2$, we say that it *selects* $V_i[r]$. Note that their lists of permitted positions mean that $P_1[0], \dots, P_k[0]$ must occur on the path in order, and observe also that every combination of starting positions for these paths can be achieved by linking the paths $P_1, \dots, P_k$ with vertices from Q. The edge-gadgets allow some vertices to be exchanged between each pair of paths P_i and P_j if the vertices selected by $P_i[0]$ and $P_j[0]$ are adjacent, and we use this, together with conditions imposed by the lists of permitted positions for $P_i[2n^2-1]$ and $P_j[2n^2-1]$, to ensure that the vertices selected by each pair of paths P_i and P_j are adjacent.

We say that a Hamilton path *swaps* in $G(e_r)$ (an edge-gadget containing edges between P_i and P_j) if the segment of the Hamilton path between $P_i[4r]$ and $P_i[4r+3]$ is

$$P_i[4r]P_j[4r+1]P_j[4r+2]P_i[4r+2]P_i[4r+3],$$

and that between $P_j[4r]$ and $P_j[4r+3]$ is

$$P_j[4r]P_i[4r+1]P_j[4r+3],$$

as illustrated in Figure 2. Observe that, if $i < j$, swapping in $G(e_r)$ increases the number of vertices lying between $P_i[0]$ and $P_i[2n^2-1]$ on the Hamilton path by one, and correspondingly decreases by one the number of vertices lying between $P_j[0]$ and $P_j[2n^2-1]$. Note that a partial swap, in which P_i gains vertices from P_j but P_j does not gain any vertex from P_i (or vice versa), is not possible in any Hamilton path, as such a path could not include $P_i[4r+1]$ (or $P_j[4r+1]$ and $P_j[4r+2]$).

Clearly every vertex of P_i that is not involved in a swap must lie between $P_i[0]$ and $P_i[2n^2-1]$ on any Hamilton path. The range of permitted positions for each vertex $P_i[l]$ (for $1 \leq i \leq k$, $1 \leq l \leq 2n^2-2$) means that, provided P_i is not involved in more than $k-1$ swaps, $P_i[l]$ will appear in a position from its list, no matter where the swaps involving P_i take place. Note also that, if $P_i[l]$ is not involved in a swap, it can only have a position from $\{(i-1)n^3 + 2rn^2 + l + \beta : -(k-1) \leq \beta \leq k-1\}$ if $P_i[0]$ selects the vertex $V_i[r]$.

In the following two propositions, we demonstrate some useful properties of this construction.

Proposition 3.2. *A valid Hamilton path can only swap in the edge-gadget $G(V_i[r]V_j[s])$ if P_i and P_j select vertices $V_i[r]$ and $V_j[s]$ respectively.*



Figure 2: A swap in $G(e_r)$

Proof. Suppose that the vertices of $G(V_i[r]V_j[s])$ are $P_i[l], \dots, P_i[l+3]$ and $P_j[l], \dots, P_j[l+3]$. Recall that

$$L_{P_i[l+1]} = L(i, l+1) \cup \{(j-1)n^3 + 2sn^2 + (l+1) + \beta : -(k-1) \leq \beta \leq k-1\}$$

and $L_{P_j[l]} = L(j, l)$. If we swap in $G(V_i[r]V_j[s])$, then $P_i[l+1]$ comes immediately after $P_j[l]$ on the Hamilton path, but if this is not possible if $P_i[l+1]$ takes a position from $L(i, l+1)$, as every element of $L(j, l)$ is strictly larger than every element of $L(i, l+1)$. Thus $P_i[l+1]$ must have a position in $\{(j-1)n^3 + 2sn^2 + (l+1) + \beta : -(k-1) \leq \beta \leq k-1\}$, implying that $P_j[l]$ has a position from the list $\{(j-1)n^3 + 2sn^2 + l + \beta : -(k-1) \leq \beta \leq k-1\}$, which is only possible if $P_j[0]$ selects $V_j[s]$.

Similarly, we see that $P_j[l+1]$ must take a position from $\{(i-1)n^3 + 2rn^2 + (l+1) + \beta : -(k-1) \leq \beta \leq k-1\}$, which implies that P_i selects $V_i[r]$. □

Proposition 3.3. *Any valid Hamilton path in H must involve exactly one swap between each pair of paths from $\{P_1, \dots, P_k\}$.*

Proof. First note that, by Proposition 3.2, there can be at most one swap between any pair of paths (since at most one edge between two given classes has both its endpoints selected). Now, the list of possible positions for $P_1[2n^2-1]$ means we must gain $k-1$ additional vertices on P_1, and this can only be achieved if there are swaps in $k-1$ gadgets involving P_1, implying that there is exactly one swap between P_1 and each other path $P_2, \ldots, P_k$.

Considering P_2, we see that it loses one vertex to P_1, so to achieve its target (a net gain of $k-3$ vertices) it must be involved in at least $k-2$ further swaps. This is the maximum possible, so again implies that in total there is exactly one swap involving P_2 and each other path $P_3, \ldots, P_k$. Continuing in this way we see that we must indeed have exactly one swap between each pair of paths from $\{P_1, \ldots, P_k\}$. □

We now use these facts to demonstrate that $(H, \mathcal{L})$ is a yes-instance for List Hamilton Path if and only if G is a yes-instance for Multicolour Clique.

Lemma 3.4. *If G is a yes-instance for* Multicolour Clique, *then $(H, \mathcal{L})$ is a yes-instance for* List Hamilton Path.

Proof. Suppose we have a multicolour clique that contains $V_i[r_i]$ for $1 \leq i \leq k$. We claim we can then find a Hamilton path in H in which every vertex has a position from its permitted list. In such a path, we give $P_i[0]$ position $(i-1)n^3 + 2r_i n^2$, using Q to link up $P_1, \ldots, P_k$ in such a way that each of these paths has the desired starting position. $P_i[0]$ occurs on the path immediately after some vertex $Q_i[\alpha]$ (where, if each path P_j for $j < i$ has a net gain of β_j vertices due to swaps, $\alpha = 2r_i n^2 - 1 - \sum_{j=1}^{i-1} \beta_j$), and $P_i[2n^2-1]$ is followed by $Q_i[\alpha+1]$, so every vertex of Q is included on the path.

We know that, for each pair (i, j) of vertex classes, there exists an edge e_{ij} between $V_i[r_i]$ and $V_j[r_j]$. Thus, for each $1 \leq i < j \leq k$, there exists a gadget $G(e_{ij})$ in which we are able to swap. If we do indeed swap in every such gadget, this gives exactly one swap between each pair of paths, so each path P_i has a net gain of $k+1-2i$ vertices and $P_i[2n^2-1]$ has position $(i-1)n^3 + 2r_i n^2 + (2n^2-1) + k + 1 - 2i \in L_{P_i[2n^2-1]}$. Each path is involved in exactly $k-1$ swaps, so all the internal vertices of $P_1, \ldots P_k$ that are not involved in swaps also receive positions from their lists.

This path then includes every vertex in H, and gives each vertex a position from its permitted list, so $(H, \mathcal{L})$ is indeed a yes-instance for List Hamilton Path. □

Lemma 3.5. *If $(H, \mathcal{L})$ is a yes-instance for* List Hamilton Path, *then G is a yes-instance for* Multicolour Clique.

Proof. Suppose there exists a valid Hamilton path in H. For $1 \leq i \leq k$, let v_i be the vertex in V_i that is selected by the position of $P_i[0]$ in this path. We claim that the set $\{v_1, \ldots, v_k\}$, which clearly contains one vertex from each colour class, induces a clique in G.

We know, by Proposition 3.3, that in any valid Hamilton path we must have exactly one swap between each pair of paths. But, by Proposition 3.2, a swap can only occur between P_i and P_j in the gadget $G(uw)$ if u and w are selected by the positions of $P_i[0]$ and $P_j[0]$ respectively. Thus a swap can only occur between paths P_i and P_j if there exists a gadget $G(v_iv_j)$, which – by construction – is only the case if there exists an edge v_iv_j in G. As this holds for every pair (v_i, v_j) with $1 \leq i < j \leq k$, we see that $\{v_1, \ldots, v_k\}$ are pairwise adjacent in G, giving the required multicolour clique. □

Finally, we need to check that H bounded pathwidth.

Lemma 3.6. *H has pathwidth at most $5k$.*

Proof. We construct a path decomposition of H, indexed by a path T with $|T| = |Q| + 2n^2 - 5$. Every bag of the decomposition contains the vertices $V_{end} = \{P_j[0], P_j[2n^2 - 1] : 1 \leq j \leq k\}$. In addition, for $1 \leq i \leq |Q| - 1$, the bag indexed by the i^{th} node of T contains the i^{th} and $(i+1)^{th}$ vertices of Q, while, for $|Q| \leq i \leq |Q| + 2n^2 - 5$, the bag indexed by the i^{th} node of T contains all vertices $\{P_j[i-|Q|+1], P_j[i-|Q|+2], P_j[i-|Q|+3] : 1 \leq j \leq k\}$. Note that every bag contains at most $5k$ vertices.

It is immediate from this construction that, for any vertex $v \in V(H)$, the nodes indexing bags that contain v induce a subpath of T. So it remains to show that, for every edge $uv \in E(H)$, there exists some bag of the decomposition that contains both u and v. For all edges within Q this is clearly true. Note that, for $1 \leq l \leq 2n^2 - 2$, any vertex $P_i[l]$ is only adjacent to vertices $P_j[l']$ where $|l - l'| \leq 2$, and so any edge between internal vertices of the paths $P_1, \ldots, P_k$ must have both its endpoints in some bag of the decomposition. All remaining edges are then incident with some $v \in V_{end}$, but V_{end} is contained in every bag, and so the condition is also satisfied for these edges.

Hence we have a path decomposition of H of width at most $5k$. □

We are now ready to prove the main theorem of this section.

Proof of Theorem 3.1. It follows immediately from Lemmas 3.4 and 3.5 that $(H, \mathcal{L})$ is a yes-instance for List Hamilton Path if and only if G is a yes-instance for Multicolour Clique. H has order polynomial in $|G|$, and can clearly be computed from G in polynomial time. Moreover, by Lemma 3.6 we know that the pathwidth of H depends only on the parameter k, the number of colours used in G. This completes the reduction to show that List Hamilton Path, parameterised by treewidth, is W[1]-hard. □

4 Conclusions and Open Problems

We have proved that List Edge Chromatic Number and List Total Chromatic Number are fixed parameter tractable, parameterised by treewidth, although the List Edge Colouring and List Total Colouring problems are NP-hard on graphs of treewidth at most two. Thus, as for vertex colouring, it is computationally easier to calculate list edge or total chromatic number of a graph than to determine whether a given set of lists admits a proper colouring of the graph.

We also demonstrated that List Hamilton Path is W[1]-hard, even when parameterised by pathwidth, giving another example of a problem that solvable in linear time on graphs of bounded treewidth but has a W[1]-hard list version.

It would be interesting to investigate whether there are further problems that are fixed parameter tractable, parameterised by treewidth, but have a W[1]-hard list version.

References

[1] Noga Alon, *Restricted colorings of graphs*, Surveys in Combinatorics, 1993, Proceedings, 14th British Combinatorial Conference, (K. Walker, ed.), London Mathematical Society Lecture Note Series, vol. 187, Cambridge University Press, 1993, pp. 1–33.

[2] Stefan Arnborg and Andrzej Proskurowski, *Linear Time Algorithms for NP-Hard Problems Restricted to Partial k-Trees*, Discrete Applied Mathematics **23** (1989), 11–24.

[3] M. Behzad, *Graphs and Their Chromatic Numbers*, Ph.D. Thesis, Michigan State University, 1965.

[4] Hans L. Bodlaender, *A Linear-Time Algorithm for Finding Tree-Decompositions of Small Treewidth*, SIAM Journal on Computing **25** (1996), no. 6, 1305–1317.

[5] B. Bollobás and A. Harris, *List-colourings of graphs*, Graphs and Combinatorics, **1** (1985), 115–127.

[6] Richard B. Borie, R. Gary Parker and Craig A. Tovey, *Automatic Generation of Linear-Time Algorithms from Predicate Calculus Descriptions of Problems on Recursively Constructed Graph Families*, Algorithmica **7** (1992), 555–581.

[7] B. Courcelle, *The monadic second-order logic of graphs I: recognizable sets of finite graphs*, Information and Computation **85** (1990), 12–75.

[8] Gruia Călinescu, Cristina G. Fernandes and Bruce Reed, *Multicuts in unweighted graphs and digraphs with bounded degree and bounded tree-width*, Journal of Algorithms **48** (2003), 333–359.

[9] R. G. Downey and M. R. Fellows, *Parameterized Complexity*, Springer-Verlag, 1999.

[10] M. Fellows, D. Hermelin, F. Rosamond and S. Vialette, *On the fixed-parameter intractability and tractability of multiple-interval graph properties*, Theoretical Computer Science **410** (2009), 53–61.

[11] Michael R. Fellows, Fedor V. Fomin, Daniel Lokshtanov, Frances Rosamond, Saket Saurabh, Stefan Szeider and Carsten Thomassen, *On the complexity of some colorful problems parameterized by treewidth*, Information and Computation **209** (2011), 143–153.

[12] F. Galvin, *The List Chromatic Index of a Bipartite Multigraph*, Journal of Combinatorial Theory, Series B **63** (1995), no. 1, 153–158.

[13] M. R. Garey, D. S. Johnson and R. Endre Tarjan, *The Planar Hamilton Circuit Problem is NP-Complete*, SIAM Journal on Computing **5** (1976), 704–714.

[14] Roland Häggkvist and Amanda Chetwynd, *Some upper bounds on the total and list chromatic numbers of multigraphs*, Journal of Graph Theory **16** (1992), no. 5, 503–516.

[15] Ian Holyer, *The NP-Completeness of Edge-Coloring*, SIAM Journal on Computing **10** (1981), no. 4, 718–720.

[16] Shuji Isobe, Xiao Zhou and Takao Nishizeki, *Total colourings of degenerate graphs*, Combinatorica **27** (2007), no. 2, 167–182.

[17] Klaus Jansen and Petra Scheffler, *Generalized coloring for tree-like graphs*, Discrete Applied Mathematics **75** (1997) 135–155.

[18] Jeff Kahn, *Asymptotics of the Chromatic Index for Multigraphs*, Journal of Combinatorial Theory, Series B **68** (1996), no. 2, 233–254.

[19] M. S. Krishnamoorthy, *An NP-hard problem in bipartite graphs*, SIGACT News **7** (1975), 26–26.

[20] Daniel Leven and Zvi Galil, *NP completeness of finding the chromatic index of regular graphs*, Journal of Algorithms **4** (1983), no. 1, 35–44.

[21] Dániel Marx, *NP-Completeness of List Coloring and Precoloring Extension on the Edges of Planar Graphs*, Journal of Graph Theory **49** (2005), 313–324.

[22] Colin J.H. McDiarmid and Abdón Sánchez-Arroyo, *Total colouring regular bipartite graphs is NP-hard*, Discrete Mathematics **124** (1994), no. 1–3, 155–162.

[23] R. Niedermeier, *Invitation to Fixed Parameter Algorithms*, Oxford University Press, 2006.

[24] D. Seese, *Tree-partite graphs and the complexity of algorithms*, Fundamentals of Computation Theory (Lothar Budach, ed.), Lecture Notes in Computer Science, vol. 199, Springer Berlin / Heidelberg 1985, pp. 412–421.

[25] V. G. Vizing, *On an estimate of the chromatic class of a p-graph. (Russian)*, Diskret. Analiz **3** (1964), 25–30.

[26] V. G. Vizing, *Some unsovled problems in graph theory (Russian)*, Uspekhi Math. Nauk. **23** (1968), 117–134.

[27] Xiao Zhou, Shin-ichi Nakano and Takao Nishizeki, *Edge-Colouring Partial k-Trees*, Journal of Algorithms **21** (1996), 598–617.

[28] Xiao Zhou, Yuki Matsuo and Takao Nishizeki, *List total colorings of series-parallel graphs*, Journal of Discrete Algorithms **3** (2005), 47–60.